



Pool of Italian Cancer Registries - 1 January 2010

MELANOMA DELLA COROIDE

CHOROIDAL MELANOMA

(ICD-10 C69 + ICD-O-3 M8720-M8790)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	475	614	769	531	279	625
% →	14%	19%	23%	16%	8%	19%
PROPORTION PER 100 000 →	0.9	1.1	1.3	1.0	0.5	1.1


MALE 54% FEMALE 46%

COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	1	6	20	28	8
NORTH EAST	1	9	19	26	8
CENTRE	0	3	18	22	5
SOUTH	0	4	15	25	5
POOL	1	6	18	26	7
FEMALE					
NORTH WEST	1	4	12	16	6
NORTH EAST	1	6	13	27	7
CENTRE	2	9	9	11	6
SOUTH	1	2	7	4	2
POOL	1	4	11	16	5
BOTH SEXES					
NORTH WEST	1	5	16	20	7
NORTH EAST	1	7	16	26	8
CENTRE	1	6	13	15	6
SOUTH	1	3	10	12	4
POOL	1	5	14	20	6

COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009


AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	33%	17%	0 - 44	>25	>25
45 - 59	21%	12%	45 - 59	>25	>25
60 - 74	11%	8%	60 - 74	>25	>25
75+	7%	6%	75+	>25	>25

2 587 347 CANCER SURVIVORS

3 292 CHOROIDAL MELANOMA CANCER SURVIVORS

ALREADY CURED at 1 JANUARY 2010

0 (0%)



COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

ICD-10 C69 + ICD-O-3 M8720-M8790 CHOROIDAL MELANOMA