




Nasopharynx cancer (ICD-10 = C11)

Tumore della rinofaringe


Observed and relative survival (%) by sex and age. Data from the Pool of Italian Cancer Registries follow-up 31.12.2003

	15-44		45-54		55-64		65-74		75+		ALL	
	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel
♂ MEN (n)	(68)		(93)		(100)		(85)		(42)		(388)	
1 year	93	93	84	84	81	82	74	76	69	75	81	83
3 years	82	83	58	59	53	55	47	51	29	37	55	59
5 years	71	71	52	53	42	44	30	36	13	20	44	48
95% CI	(60-81)	(60-82)	(41-62)	(42-63)	(32-52)	(34-55)	(21-40)	(24-47)	(3-24)	(3-36)	(39-49)	(43-54)
♀ WOMEN (n)	(29)		(28)		(27)		(27)		(30)		(141)	
1 year	97	97	100	100	78	78	74	75	47	49	79	80
3 years	83	83	75	75	63	64	56	58	20	24	59	62
5 years	79	80	64	65	59	60	44	47	10	14	51	56
95% CI	(65-94)	(65-94)	(46-82)	(47-83)	(41-78)	(41-79)	(25-63)	(27-68)	(-1-21)	(-1-29)	(43-59)	(46-65)
ALL (n)	(97)		(121)		(127)		(112)		(72)		(529)	
1 year	94	94	88	88	80	88	74	76	60	64	80	82
3 years	83	83	62	63	55	57	49	52	25	31	56	60
5 years	73	74	54	55	45	48	34	39	12	18	46	50
95% CI	(64-82)	(65-82)	(46-63)	(46-64)	(37-54)	(39-57)	(25-43)	(29-49)	(4-20)	(6-29)	(41-50)	(45-55)


Five-year age-standardised relative survival



Relative survival by Italian Areas



Trend of relative survival Pool of some Italian Registries



→ Ulteriori dati sono disponibili presso: www.registri-tumori.it