




Hipopharynx cancer (ICD-10 = C12-13)

Tumore dell'ipofaringe


Observed and relative survival (%) by sex and age. Data from the Pool of Italian Cancer Registries follow-up 31.12.2003

	15-44		45-54		55-64		65-74		75+		ALL	
	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel
♂ MEN (n)	(38)		(185)		(282)		(225)		(78)		(808)	
1 year	58	58	84	84	75	76	67	69	56	62	72	74
3 years	37	37	51	52	40	42	35	38	17	21	39	41
5 years	34	35	38	39	32	34	26	30	10	15	29	33
95% CI	(19-49)	(19-50)	(31-45)	(32-46)	(26-37)	(28-39)	(20-31)	(23-37)	(3-16)	(4-26)	(26-33)	(29-36)
♀ WOMEN (n)	(7)		(12)		(34)		(26)		(23)		(102)	
1 year	71	72	83	83	68	68	54	55	48	51	62	63
3 years	29	29	67	67	35	36	42	44	26	32	38	41
5 years	14	14	50	51	27	27	34	37	17	23	28	32
95% CI	(-12-40)	(-12-40)	(22-78)	(22-79)	(12-41)	(12-42)	(16-53)	(17-57)	(2-33)	(0-47)	(20-37)	(22-42)
ALL (n)	(45)		(197)		(316)		(251)		(101)		(910)	
1 year	60	60	84	84	74	38	66	68	55	59	71	73
3 years	36	36	52	53	40	41	36	39	19	24	39	41
5 years	31	31	39	40	31	33	27	31	12	18	29	33
95% CI	(18-45)	(18-45)	(32-46)	(33-47)	(26-36)	(27-38)	(21-32)	(24-37)	(5-18)	(8-28)	(26-32)	(29-36)


Five-year age-standardised relative survival



Relative survival by Italian Areas



Trend of relative survival Pool of some Italian Registries



➔ Ulteriori dati sono disponibili presso: www.registri-tumori.it