2 587 347 CANCER SURVINGE



Pool of Italian Cancer Registries - 1 January 2010

OROFARINGE OROPHARYNX

(ICD-10 C09-10)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	1 578	1 300	1 386	781	512	680
% →	25%	21%	22%	13%	8%	11%
PROPORTION → PER 100 000	3	3	3	2	1	1

MALE 77% FEMALE 23%

COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE


(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	1	23	72	90	26
NORTH EAST	1	25	82	70	25
CENTRE	0	11	40	38	12
SOUTH	1	12	27	32	9
POOL	1	19	60	63	19
FEMALE					
NORTH WEST	0	7	20	13	7
NORTH EAST	1	10	20	17	8
CENTRE	1	6	5	5	3
SOUTH	0	5	3	5	2
POOL	0	7	14	11	5
BOTH SEXES					
NORTH WEST	0	15	44	40	16
NORTH EAST	1	17	49	36	16
CENTRE	0	8	22	18	7
SOUTH	0	8	14	15	5
POOL	0	13	36	30	12


ALREADY CURED at 1 JANUARY 2010

6 237 **OROPHARYNX CANCER SURVIVORS**

489 (24%)



COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

	CURE FRACTION %		
AGE AT DIAGNOSIS	MALE	FEMALE	
0 - 44	32%	33%	
45 - 59	24%	31%	
60 - 74	15%	29%	
75 +	12%	27%	

	TIME TO CURE YEARS				
AGE AT DIAGNOSIS	MALE	FEMALE			
0 - 44	11	11			
45 - 59	12	11			
60 - 74	13	11			
75+	13	12			

COMPLETE PREVALENCE

Overall number (or proportion) of cancer survivors

TIME TO CURE

Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS

Patients who have survived longer than time to cure.

CURE FRACTION

Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.