




Pool of Italian Cancer Registries - 1 January 2010

# RINOFARINGE NASOPHARYNX

(ICD-10 C11)

## COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	1 025	940	1 139	842	743	3 429
% →	13%	12%	14%	10%	9%	42%
PROPORTION PER 100 000 →	2	2	2	1	1	6
MALE 51%			FEMALE 49%			




## COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)


AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
<b>MALE</b>					
NORTH WEST	4	18	35	25	15
NORTH EAST	3	19	25	40	13
CENTRE	2	13	14	21	8
SOUTH	6	31	43	23	18
POOL	4	22	33	29	15
<b>FEMALE</b>					
NORTH WEST	3	13	22	13	10
NORTH EAST	2	12	24	28	11
CENTRE	4	17	17	28	12
SOUTH	5	28	49	23	18
POOL	3	18	30	22	13
<b>BOTH SEXES</b>					
NORTH WEST	4	15	28	14	12
NORTH EAST	2	15	24	33	12
CENTRE	3	15	16	25	10
SOUTH	6	28	46	20	17
POOL	4	19	31	23	14

ALREADY CURED at 1 JANUARY 2010

3 939 (49%)



## COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



**COMPLETE PREVALENCE**  
Overall number (or proportion) of cancer survivors.

**TIME TO CURE**  
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

**ALREADY CURED SURVIVORS**  
Patients who have survived longer than time to cure.

**CURE FRACTION**  
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

## CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	50%	54%	0 - 44	12	12
45 - 59	29%	38%	45 - 59	14	17
60 - 74	10%	21%	60 - 74	16	22
75+	4%	14%	75+	17	>25