



Pool of Italian Cancer Registries - 1 January 2010

STOMACO

STOMACH

(ICD-10 C16)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	14 396	12 347	15 306	10 975	7 062	14 331
% →	19%	17%	21%	15%	9%	19%
PROPORTION PER 100 000 →	25	21	26	19	12	25


MALE 55% **FEMALE 45%**

COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)


AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	3	81	374	855	159
NORTH EAST	5	81	417	1 094	178
CENTRE	7	116	460	1 159	204
SOUTH	4	60	235	492	83
POOL	4	77	354	856	144
FEMALE					
NORTH WEST	4	50	218	454	118
NORTH EAST	5	57	254	668	147
CENTRE	5	82	302	688	157
SOUTH	4	45	133	263	59
POOL	4	53	211	495	111
BOTH SEXES					
NORTH WEST	4	65	290	598	137
NORTH EAST	5	69	331	825	162
CENTRE	6	99	377	871	180
SOUTH	4	52	181	352	70
POOL	4	65	278	630	127

COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	40%	42%	0 - 44	8	8
45 - 59	29%	32%	45 - 59	9	9
60 - 74	18%	22%	60 - 74	10	10
75+	13%	17%	75+	11	11



NORTH WEST CANCER REGISTRY AREAS
NORTH EAST CANCER REGISTRY AREAS
CENTRE CANCER REGISTRY AREAS
SOUTH CANCER REGISTRY AREAS

COMPLETE PREVALENCE
 Overall number (or proportion) of cancer survivors.

TIME TO CURE
 Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
 Patients who have survived longer than time to cure.

CURE FRACTION
 Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

ICD-10 C16 STOMACH