

Pool of Italian Cancer Registries - 1 January 2010

# FEGATO

## LIVER

(ICD-10 C22)

### COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS



YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	10 693	7 245	3 841	1 263	421	208
% →	45%	31%	16%	5%	2%	1%
PROPORTION PER 100 000 →	19	13	7	2	1	0
MALE 70%			FEMALE 30%			

### COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
<b>MALE</b>					
NORTH WEST	5	67	193	261	74
NORTH EAST	4	48	200	266	67
CENTRE	1	34	125	162	42
SOUTH	1	40	194	199	51
POOL	3	49	191	237	62
<b>FEMALE</b>					
NORTH WEST	2	12	49	100	27
NORTH EAST	2	10	43	101	24
CENTRE	3	7	33	68	17
SOUTH	3	9	84	82	24
POOL	2	10	56	94	24
<b>BOTH SEXES</b>					
NORTH WEST	4	39	116	158	49
NORTH EAST	3	29	118	161	45
CENTRE	2	20	77	105	29
SOUTH	2	24	136	128	37
POOL	3	29	119	147	42

### COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



ICD-10 C22 LIVER



**COMPLETE PREVALENCE**  
Overall number (or proportion) of cancer survivors.

**TIME TO CURE**  
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

**ALREADY CURED SURVIVORS**  
Patients who have survived longer than time to cure.

**CURE FRACTION**  
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

### CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	6%	13%	0 - 44	22	20
45 - 59	3%	6%	45 - 59	24	23
60 - 74	1%	2%	60 - 74	25	25
75+	1%	1%	75+	>25	>25