

Pool of Italian Cancer Registries - 1 January 2010


TESSUTO CONNETTIVO

CONNECTIVE TISSUE

(ICD-10 C47,49)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	2 971	3 570	4 253	2 979	2 017	5 646
% →	14%	17%	20%	14%	9%	26%
PROPORTION PER 100 000 →	5	7	8	5	3	10
	MALE 53%			FEMALE 47%		




COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)


AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	16	39	88	134	44
NORTH EAST	19	53	88	135	47
CENTRE	20	45	62	129	41
SOUTH	18	36	64	86	33
POOL	18	43	79	120	41
FEMALE					
NORTH WEST	15	43	60	100	41
NORTH EAST	16	38	67	81	38
CENTRE	11	31	38	61	26
SOUTH	15	39	49	41	28
POOL	15	39	58	75	35
BOTH SEXES					
NORTH WEST	16	41	73	112	42
NORTH EAST	18	46	77	100	42
CENTRE	16	38	50	87	33
SOUTH	17	38	56	58	30
POOL	17	41	68	92	38

ALREADY CURED at 1 JANUARY 2010

11 085 (52%)



COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	69%	67%	0 - 44	7	7
45 - 59	53%	51%	45 - 59	9	9
60 - 74	40%	39%	60 - 74	11	11
75+	33%	33%	75+	12	11