



Pool of Italian Cancer Registries - 1 January 2010


MAMMELLA FEMMINILE

FEMALE BREAST

(ICD-10 C50)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	86 260	113 597	151 262	99 546	62 718	67 989
% →	15%	20%	26%	17%	11%	12%
PROPORTION PER 100 000 →	297	390	525	348	220	240



COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE



(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
FEMALE					
NORTH WEST	201	2 491	5 302	6 297	2 495
NORTH EAST	188	2 514	5 325	6 030	2 332
CENTRE	159	2 101	3 967	4 490	1 749
SOUTH	146	1 725	3 431	3 957	1 356
POOL	175	2 223	4 676	5 455	2 019

ALREADY CURED at 1 JANUARY 2010

90 331 (16%)

COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



ICD-10 C50 FEMALE BREAST

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %
	FEMALE
0 - 44	46%
45 - 59	64%
60 - 74	52%
75+	47%

AGE AT DIAGNOSIS	TIME TO CURE YEARS
	FEMALE
0 - 44	>25
45 - 59	15
60 - 74	20
75+	17

COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.