



Pool of Italian Cancer Registries - 1 January 2010


CERVICE UTERINA

CERVIX UTERI

(ICD-10 C53)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	4 253	4 933	7 747	6 852	6 457	30 051
% →	7%	8%	13%	11%	11%	50%
PROPORTION PER 100 000 →	14	17	25	24	21	102




ALREADY CURED at 1 JANUARY 2010
44 717 (74%)


COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
FEMALE					
NORTH WEST	37	238	434	703	247
NORTH EAST	32	215	384	595	205
CENTRE	26	211	366	586	192
SOUTH	18	180	418	515	163
POOL	28	209	407	607	202



COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	FEMALE			FEMALE	
0 - 44	76%		0 - 44	6	
45 - 59	62%		45 - 59	8	
60 - 74	41%		60 - 74	10	
75+	29%		75+	11	