



Pool of Italian Cancer Registries - 1 January 2010


OVAIO

OVARY

(ICD-10 C56)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	7 083	7 018	7 926	6 241	4 290	10 023
% →	17%	16%	19%	15%	10%	24%
PROPORTION PER 100 000 →	24	23	26	21	14	34




COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)


AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
FEMALE					
NORTH WEST	25	178	366	330	164
NORTH EAST	24	190	332	322	151
CENTRE	35	163	305	307	141
SOUTH	27	171	283	229	116
POOL	26	179	326	299	142

ALREADY CURED at 1 JANUARY 2010

17 226 (40%)



COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



FEMALE
ALREADY CURED

NORTH WEST CANCER REGISTRY AREAS
NORTH EAST CANCER REGISTRY AREAS
CENTRE CANCER REGISTRY AREAS
SOUTH CANCER REGISTRY AREAS

ICD-10 C56 OVARY

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	FEMALE			FEMALE	
0 - 44	59%		0 - 44	9	
45 - 59	38%		45 - 59	11	
60 - 74	15%		60 - 74	14	
75+	7%		75+	16	

COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.