



Pool of Italian Cancer Registries - 1 January 2010


PROSTATA

PROSTATE

(ICD-10 C61)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS

YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	66 485	93 773	95 239	30 255	8 131	1 740
% →	22%	32%	32%	10%	3%	1%
PROPORTION PER 100 000 →	245	352	364	114	32	6



COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE



(PROPORTION PER 100 000)

AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	1	306	3 821	7 929	1 428
NORTH EAST	2	313	3 933	8 256	1 395
CENTRE	0	165	2 582	6 447	1 015
SOUTH	3	128	1 806	4 286	588
POOL	2	243	3 184	6 881	1 114

ALREADY CURED at 1 JANUARY 2010

320 (0.1%)

COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



- NORTH WEST** CANCER REGISTRY AREAS
- NORTH EAST** CANCER REGISTRY AREAS
- CENTRE** CANCER REGISTRY AREAS
- SOUTH** CANCER REGISTRY AREAS

ICD-10 C61 PROSTATE

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %	
	MALE	
0 - 44	72%	
45 - 59	59%	
60 - 74	42%	
75+	32%	

AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	
0 - 44	10	
45 - 59	17	
60 - 74	25	
75+	>25	

COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.