



Pool of Italian Cancer Registries - 1 January 2010

RENE E ALTRI ORGANI URINARI KIDNEY AND OTHER URINARY ORGANS

(ICD-10 C64-66,68)

COMPLETE PREVALENCE BY YEARS SINCE DIAGNOSIS



YEARS →	≤2	(2 - 5]	(5 - 10]	(10 - 15]	(15 - 20]	> 20
No. →	17 449	20 315	24 252	15 102	8 652	11 870
% →	18%	21%	25%	15%	9%	12%
PROPORTION PER 100 000 →	31	36	44	27	16	21
MALE 62%			FEMALE 38%			

COMPLETE PREVALENCE BY SEX, MACRO-AREA, AND AGE

(PROPORTION PER 100 000)


AGE CLASS →	0-44	45-59	60-74	75+	ALL AGES
MALE					
NORTH WEST	20	198	642	1 177	268
NORTH EAST	23	211	741	1 296	286
CENTRE	22	189	611	954	234
SOUTH	21	132	370	486	126
POOL	22	180	595	1 002	225
FEMALE					
NORTH WEST	22	89	269	483	148
NORTH EAST	23	117	332	570	169
CENTRE	23	88	271	388	125
SOUTH	21	72	170	202	73
POOL	22	93	262	433	129
BOTH SEXES					
NORTH WEST	21	142	441	731	205
NORTH EAST	23	164	526	835	226
CENTRE	23	137	433	608	178
SOUTH	21	101	264	313	99
POOL	22	136	418	644	175

COMPLETE PREVALENCE AND PROPORTION OF ALREADY CURED SURVIVORS BY SEX



ALREADY CURED at 1 JANUARY 2010

8 965 (9%)



COMPLETE PREVALENCE
Overall number (or proportion) of cancer survivors.

TIME TO CURE
Time span after cancer diagnosis necessary to eliminate excess mortality due to cancer. It is measured as the time necessary to reach a 5-year conditional relative survival (that is the probability to survive additional five years) >95%.

ALREADY CURED SURVIVORS
Patients who have survived longer than time to cure.

CURE FRACTION
Proportion of cancer patients who are expected to reach the same death rates of the general population and will not die as a result of their cancer.

CURE FRACTION AND TIME TO CURE BY AGE AT DIAGNOSIS AND SEX FOR CANCER PATIENTS DIAGNOSED IN ITALY IN 1985-2009

AGE AT DIAGNOSIS	CURE FRACTION %		AGE AT DIAGNOSIS	TIME TO CURE YEARS	
	MALE	FEMALE		MALE	FEMALE
0 - 44	59%	71%	0 - 44	11	6
45 - 59	44%	56%	45 - 59	21	11
60 - 74	27%	37%	60 - 74	>25	19
75+	18%	26%	75+	>25	25