


NASOPHARYNX (ICD-IX 147)

Pool of Italian Registries

REGISTRY	Number of cases		
	Men	Women	Total
Torino	24	5	29
Genova	24	5	29
Varese	23	10	33
Veneto	44	21	65
Parma	9	2	11
Modena	19	4	23
Ferrara	11	6	17
Romagna	13	7	20
Macerata	2	1	3
Firenze-Prato	25	15	40
Umbria	14	6	20
Latina	8	0	8
Sassari	5	0	5
Ragusa	8	5	13
Italy	229	87	316



Observed and relative survival (%) by sex and age classes

	15-44		45-54		55-64		65-74		75+		ALL	
	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel	obs	rel
Men												
1 year	94	94	79	79	76	77	66	67	37	40	73	75
3 years	73	74	58	59	57	59	43	46	11	14	52	55
5 years	64	65	36	37	45	49	26	30	0	0	38	42
95% CI	(47-78)	(47-79)	(23-51)	(23-52)	(34-57)	(37-61)	(16-38)	(19-44)	-	-	(31-44)	(35-49)
Women												
1 year	95	96	85	85	86	87	72	73	50	53	80	81
3 years	91	91	54	54	50	51	50	52	50	59	61	63
5 years	71	72	45	45	41	42	33	36	42	57	48	51
95% CI	(50-86)	(50-87)	(22-70)	(22-71)	(23-61)	(24-63)	(16-56)	(18-61)	(19-68)	(26-93)	(37-58)	(40-62)
Total												
1 year	95	95	80	81	79	80	67	69	42	45	75	77
3 years	80	81	57	58	55	57	44	48	26	33	54	57
5 years	67	68	38	39	44	47	27	31	18	28	40	44
95% CI	(54-78)	(54-79)	(26-51)	(27-53)	(34-54)	(37-58)	(19-38)	(21-44)	(8-36)	(12-55)	(35-46)	(38-50)